

THE TOPOLOGY OF WEAK LENSING FIELDS

TAKAHIKO MATSUBARA

Department of Physics and Astrophysics, Nagoya University, Chikusa-ku, Nagoya 464-8602, Japan

AND

BHUVNESH JAIN

Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104

taka@a.phys.nagoya-u.ac.jp, bjain@hep.upenn.edu

Draft version February 1, 2008

ABSTRACT

The topology of weak lensing fields is studied using the 2-dimensional genus statistic. Simulated fields of the weak lensing convergence are used to focus on the effect of nonlinear gravitational evolution and to model the statistical errors expected in observational surveys. For large smoothing angles, the topology is in agreement with the predictions from linear theory. On smoothing angles smaller than $10'$, the genus curve shows the non-Gaussian signatures of gravitational clustering and differs for open and flat cold dark matter models. Forthcoming surveys with areas larger than 10 square degrees should have adequate signal-to-noise to measure the non-Gaussian shape and the Ω -dependence of the genus statistic.

Subject headings: cosmology: theory — gravitational lensing — large-scale structure of universe — methods: numerical

1. INTRODUCTION

Mapping the mass density field with weak gravitational lensing is a promising method to analyze large-scale structure in the universe. The statistical nature of the mass density field is directly predicted by theories of the early universe. For instance, most inflationary theories predict a random Gaussian initial mass density field, while structure formation by cosmic defects predicts a highly non-Gaussian field. As the density field evolves, gravitational clustering produces characteristic non-Gaussian features. Thus measuring the non-Gaussianity of the density field can be an important test of the origin and evolution of large-scale structure.

The genus statistic is known to be a useful topological descriptor of the clustering pattern of the mass density field (Gott, Melott & Dickinson 1986; Gott et al. 1989). The “genus curve”, which plots the measured genus versus the threshold for iso-density contours, has a universal shape for a random Gaussian field. Departures from this Gaussian shape reveal the mechanism of the formation of nonlinear structures. So far, the genus statistic has been applied to the distribution of galaxies, and of clusters (Moore et al. 1992; Rhoads, Gott & Postman 1994; Canavezes et al. 1998). Since these tracers are affected by biasing, it is desirable to directly use the mass density field for the genus statistic. Fortunately, recent developments in measuring weak lensing in wide area surveys suggest that this may be feasible in the near future (van Waerbeke et al. 2000; Bacon Refregier & Ellis 2000; Wittman et al. 2000; Kaiser, Wilson & Luppino 2000). As weak lensing maps of the convergence field, κ , cover larger areas with better signal-to-noise, statistical properties beyond just the second moment can be measured. It is therefore important to obtain theoretical predictions for the behavior of the genus statistic for the weak lensing field, which we present in this paper.

The linear-scale fluctuations of the mass density field

preserves the Gaussianity of the initial state. However, measuring the linear-scale fluctuations is difficult due to sample variance, unless the area covered is extremely large. In forthcoming weak lensing surveys, the convergence field is likely to contain non-linear effects of gravitational clustering. The genus curve is distorted by the non-Gaussianity induced by gravitational evolution, which precludes a simple comparison of the observed genus with analytical predictions. Since this effect is dependent on the cosmological model, it is of interest to establish whether forthcoming surveys will allow for an adequate measurement of the non-Gaussian genus curve.

We show in this paper that the effects of gravitational clustering on the genus curve can be measured from high quality data. For this purpose, we analyze the ray-tracing numerical simulation of weak lensing developed by Jain, Seljak & White (2000). The noise due to the finite number of galaxies used to probe the weak lensing field reduces the signal-to-noise on very small scales. Therefore we have to balance sample variance on large scales with discreteness noise on small scales to find the best length-scales for measuring the genus. We study this issue by simulating the random noise expected in observational surveys.

This paper is organized as follows. §2 introduces the genus statistic. In §3 the main results of our analysis from ray tracing simulations are presented. The genus curves of simulated noise-free convergence fields are shown in §3.1 and those of noisy fields in §3.2. We conclude in §4.

2. GENUS STATISTICS AND WEAK LENSING

We consider the local convergence field κ and calculate genus statistics of this field. In the weak lensing regime, $\kappa \ll 1$, an area in the lens plane is magnified by a factor $1 + 2\kappa$ compared to that in the source plane. The local convergence field can be reconstructed from the shear field (Kaiser & Squires 1993). For a distant observer, under the Born approximation, the convergence along a line of sight

can be expressed as a projection of the density perturbation δ ,

$$\kappa = \frac{3H_0^2\Omega_0}{2} \int_0^\chi \frac{d_A(\chi')d_A(\chi - \chi')}{d_A(\chi)} \frac{\delta(\chi')}{a(\chi')} d\chi', \quad (1)$$

where Ω_0 is the present value of the mass density parameter, H_0 is the Hubble's constant, $a(\chi)$ is the scale factor at the conformal lookback time χ , $d_A(\chi)$ is the comoving angular diameter distance, χ the comoving radial coordinate. If the radial distance is measured in units of H_0^{-1} , we can drop the H_0^2 in the above equation. Integration is performed along the unperturbed light path. In this case, the statistics of the convergence field can be computed from the statistics of the density field.

To quantify the topology, we find contour lines of a threshold value labelled by ν . For a Gaussian random field, the threshold ν is identified so that the field takes the value $\kappa = \nu\sigma$, where σ is the rms of the field, $\sigma^2 = \langle \kappa^2 \rangle$. The genus is defined as: (number of contours surrounding regions higher than the threshold value) - (number of contours surrounding regions lower than the threshold value) (Adler 1981; Coles 1988; Melott et al. 1989; Gott et al. 1990). The genus per unit area $G_2(\nu)$, as a function of the threshold is known as the genus curve.

The genus curve is analytically given by (Melott et al. 1989):

$$G_2(\nu) = \frac{1}{(2\pi)^{3/2}} \frac{\langle k^2 \rangle}{2} \nu e^{-\nu^2/2}, \quad (2)$$

where $\langle k^2 \rangle$ is the square of the wave number k averaged over the smoothed 2D power spectrum. Both sides of the above equation have dimensions of [steradian] $^{-2}$. If the 3D power spectrum is given by a power law with spectral index n , and a Gaussian window function $W(r) \propto e^{-r^2/(2\theta_f^2)}$ is adopted, then $\langle k^2 \rangle = (n+2)/(2\theta_f^2)$ for $n > -2$.

The above analytic results are not expected to hold for ongoing weak lensing surveys because the survey areas are not large enough to probe the linear scales. However, it is known that the shape of the genus curve of mass density fields is not strongly affected by nonlinear gravitational evolution provided we use a rescaled labeling of ν (Gott, Weinberg & Melott 1987; Weinberg, Gott & Melott 1987; Melott, Weinberg & Gott 1988). For a non-Gaussian field, the threshold is defined through the fraction of area f on the high-density side of the contour lines, which is related to the corresponding values of ν for a Gaussian field as:

$$f = \frac{1}{\sqrt{2\pi}} \int_\nu^\infty e^{-t^2/2} dt. \quad (3)$$

For a Gaussian field, this definition gives the same result as specifying the threshold directly in units of σ . For a non-Gaussian field, it compensates for the horizontal shift of the genus curve due to the shift of the probability distribution function. We adopt this prescription for setting the threshold values of κ in the results shown below.

3. NUMERICAL RESULTS

3.1. Genus without noise

We use maps of convergence fields from the ray-tracing simulations of Jain, Seljak & White (2000). The source galaxies are taken to be at $z = 1$. We use two cold dark matter models: a ‘‘tau’’ model with $\Omega_0 = 1$, and shape

parameter $\Gamma = 0.21$, and an open model with $\Omega_0 = 0.3$, $\Gamma = 0.21$. The fields we use are 166.28 arcminutes on a side for the tau model, and 235.68 arcminutes for the open model, sampled with 2048x2048 grids (see Jain, Seljak, & White 2000, for further details). Figure 1 shows a κ map for the open model.

The genus curves are calculated using the CONTOUR2D code and related routines, which were kindly provided by David Weinberg (see Gott, Melott & Dickinson (1986) and Melott (1990) for details of the methods). The key relation used is a 2-dimensional analog of the Gauss-Bonnet theorem, which expresses the genus in terms of the integral of the curvature along a contour line, $G_2 = \int C ds/2\pi$, where C is given by the inverse radius of curvature r^{-1} , and is negative or positive, depending on whether a low or high density region is enclosed. For each model, genus curves are obtained for 7-10 realizations. The error bars on the genus curves shown in Figures 2 and 3 give the 1σ deviations among those realizations, and represent the 1σ error expected for surveys of our sample area.

In the left panels of each plot in Figure 2, the genus curves per steradian of the simulated convergence fields are plotted for the open and the tau models. Each plot has a different smoothing angle, $\theta_f = 1', 2', 3', 10'$. In the plots, thin curves are the Gaussian predictions with normalizations fitted by minimizing chi-squares of the simulation data. The smoothing angle of $10'$ corresponds to a scale on which linear theory can be applied (e.g., see Fig. 3 of Jain & Seljak 1997). Within the sample variance error bars, the genus curves of the simulation data are consistent with Gaussian predictions on this linear scale.

On smaller, nonlinear scales of $\theta_f = 1', 2', 3'$, the genus curves of the convergence field show a ‘‘meatball shift’’: a dominance of the number of isolated clusters over isolated voids. This shift is consistent with topological analyses of the galaxy distributions (Gott et al. 1989, 1992), and is a common feature of cold dark matter models (Springel et al. 1998). To quantify the degree of the meatball shift, we define a meatball-shift statistic, $S = \int G_2(\nu) d\nu / \int |G_2(\nu)| d\nu$, where the integration is limited from $\nu = -2$ to $+2$. Likewise, the departure from Gaussianity can be quantified by a statistic like: $S_2 = \int |G_2(\nu) - G_{\text{gauss}}(\nu)| / \int |G_{\text{gauss}}|$. We discuss some results in the following sub-section.

The absolute values of the genus number of the high-convergence side ($\nu > 0$) is larger for the open model, and smaller for the tau model. Since our open and tau models have the same initial power spectrum, the difference in the genus number is due to the difference in the nonlinear gravitational evolution, and, more importantly, in the space-time geometry. A volume factor $V(z)$ for a unit solid angle is larger in the open model than in the tau model. Nonlinear structures also form earlier in the open model. Both these effects cause the topology statistics for the open model to have different non-Gaussian features from the tau model.

3.2. Genus with realistic noise

The weak lensing fields in observational surveys contain noise due to the finite number of source galaxies with randomly oriented intrinsic ellipticities. We build simulated noisy maps of the convergence field by drawing each

component of the intrinsic ellipticity of a galaxy from a Gaussian with variance (e.g., van Waerbeke 2000)

$$\sigma_{\text{noise}}^2 = \frac{\sigma_\epsilon^2}{2N_g}, \quad (4)$$

where σ_ϵ is the rms amplitude of the intrinsic ellipticity distribution and N_g is the mean number density of source galaxies in each cell. In the following, the number density of source galaxies is 30 per square arcminute, and their rms intrinsic ellipticity is $\sigma_\epsilon = 0.2$.

Figure 3 shows the genus measured from a κ field reconstructed from pure Gaussian noise which corresponds to a map in which galaxies are randomly distributed with random ellipticities without weak lensing signals. In fact, any white noise has the same genus curve of equation (2) regardless of its variance. The agreement with the analytic curve for a random Gaussian field (Eq. 2) tests our reconstruction scheme and genus measurement. In Figure 2 the right panels in each plot show the genus curves of simulated signal+noise maps of the convergence fields.

For the large smoothing angle $\theta_f = 10'$, the genus curves are again consistent with the Gaussian predictions. For $\theta_f = 3'$, the noise effect is small, but the meatball shift is reduced from the noise-free maps. For $\theta_f = 2'$, the meatball shift is completely erased by noise except in the vicinity of $\nu = 0$. For $\theta_f = 1'$, there are interesting features due to the noise. The relative effect of noise is stronger for the open model because the lensing signal is small [see Eq. 1]. The high-convergence region of $\nu \sim 2$ is not much affected by the noise. This is because most of the high-convergence peaks come from strongly clustered regions. Around $\nu \sim 1$, the noise contribution equals the signal. For the low-convergence region of $-\nu \sim 1-2$ the genus is dominated by noise because relatively large areas of low-convergence (which corresponds to voids) have noise troughs superimposed on them. Such complex situations make the genus curve have larger amplitude than the Gaussian at negative ν for $\theta_f = 1'$ —the opposite of the case for the noise-free maps. There is an interesting counterpart to this effect in the statistics of peaks studied by Jain, van Waerbeke (2000); they also found an increase in the number of negative peaks (relative to the Gaussian case) in the signal+noise maps.

These effects on the genus curve discussed above can

be quantified using the statistics giving the meatball shift and departure from Gaussianity, S and S_2 . For the meatball shift statistic we find that the open model shows a significant negative (sponge-like) shift for $\theta = 1'$: $S = -0.08 \pm 0.01$. It shows a positive (meatball) shift for large angles, e.g. for $\theta = 3'$, $S = 0.06 \pm 0.02$. The tau model shows a similar trend but with lower amplitude. This is in contrast to the noise-free field, for which there is a positive meatball shift which decreases in amplitude with increasing angular scale.

4. DISCUSSION

We have analyzed the genus curve of weak lensing convergence fields. On large angular scales, the genus curves are consistent with the random Gaussian prediction. On scales of a few arcminutes, nonlinear gravitational clustering produces a non-Gaussian “meatball” shift in the genus curve. Its non-Gaussianity also differentiates the open and flat cosmological model for a range of smoothing angles and threshold values. The noise dilutes the meatball shift but it does not change the genus curve to the random Gaussian one, even on the smallest scale $\theta_f = 1'$. Over the range of scales tested, $\theta_f = 1'-10'$, the open and flat cold dark matter models can be distinguished even in the presence of noise.

We have analysed simulations of survey areas of about 10 square degrees. When survey areas approach 100 square degrees, the error bars shown in Fig. 2 should reduce by about a factor of three. Thus, provided systematic errors can be kept to below the level of the random noise modeled here, the genus statistic can provide a valuable means of measuring the non-Gaussian features of the lensing signal and distinguishing cosmological models. Further work to measure other topological statistics and simulate a wider set of cosmological models would be useful.

We are very grateful to David Weinberg for providing us useful routines to calculate the 2D genus curve and for stimulating discussions. We thank an anonymous referee for suggestions that helped improve the paper. T. M. acknowledges support from JSPS Postdoctoral Fellowships for Research Abroad. B. J. acknowledges support from NASA through grants NAG5-9186 and NAG5-9220.

REFERENCES

- Adler, R. J. 1981, R. K. 1986, *The Geometry of Random Fields* (Chichester: Wiley)
- Bacon, D., Refregier, A. & Ellis, R. 2000, to appear in Proceedings of XXth Moriond Astrophysics Meeting, “Cosmological Physics with Gravitational Lensing”, eds. J.-P. Kneib, Y. Mellier, M. Moniez and J. Tran Thanh Van (astro-ph/0008248)
- Canavezes, A., Springel, V. et al. 1998, *MNRAS*, 297, 777
- Coles, P. 1988, *MNRAS*, 234, 509
- Gott, J. R., Weinberg, D. H. & Melott, A. L. 1987, *ApJ*, 319, 1
- Gott, J. R., Melott, A. L. & Dickinson, M. 1986, *ApJ*, 306, 341
- Gott, J. R. et al. 1989, *ApJ*, 340, 625
- Gott, J. R., Park, C., Juszkiewicz, R., Bies, W. E., Bennett, D. P., Bouchet, F. R. & Stebbins, A. 1990, *ApJ*, 352, 1
- Gott, J. R., Mao, S., Park, C. & Lahav, O. 1992, *ApJ*, 385, 26
- Jain, B. & Seljak, U. 1997, *ApJ*, 484, 560
- Jain, B., Seljak, U. & White, S. 2000, *ApJ*, 530, 547
- Jain, B. & van Waerbeke, L. 2000, *ApJ*, 530, L1
- Kaiser, N. & Squires, G. 1993, *ApJ*, 404, 441
- Kaiser, N., Wilson, G. & Luppino, G. A. 2000, submitted to *ApJ Letters* (astro-ph/0003338)
- Melott, A. L. 1990, *Phys. Rep.*, 193, 1
- Melott, A. L., Weinberg, D. H. & Gott, J. R. 1988, *ApJ*, 328, 50
- Melott, A. L., Cohen, A. P., Hamilton, A. J. S., Gott, J. R. & Weinberg, D. H. 1989, *ApJ*, 345, 618
- Moore, B. et al. 1992, *MNRAS*, 256, 477
- Rhoads, J. E., Gott, J. R. I. & Postman, M. 1994, *ApJ*, 421, 1
- Springel, V. et al. 1998, *MNRAS*, 298, 1169
- van Waerbeke, L. 2000, *MNRAS*, 313, 524
- van Waerbeke, L. et al. 2000, *A&A*, 358, 30
- Weinberg, D. H., Gott, J. R. & Melott, A. L. 1987, *ApJ*, 321, 2
- Wittman, D. M., Tyson, J. A., Kirkman, D., Dell’Antonio, I. & Bernstein, G. 2000, *Nature*, 405, 143




FIG. 1.— Comparison of the reconstructed and input maps of κ . The left panel shows the input κ maps for a field 3 degrees on a side, and the right panel shows the same field reconstructed from simulated ellipticity data that includes the noise due to the intrinsic ellipticities of galaxies.




FIG. 2.— Genus curves per steradian for simulated convergence fields. The four plots use smoothing angles $\theta_f = 1', 2', 3', 10'$. In each plot, fields without noise (left panels) and with noise due to the intrinsic ellipticities of galaxies (right panels) are shown. Symbols and thick lines correspond to the genus measured from the simulation data. Thin curves correspond to the prediction of random Gaussian fields with normalization fitted to the simulation data. Two cosmological models are shown: the open model (solid), and the tau model (dashed).




FIG. 3.— The genus curve of the pure Gaussian noise. This curve corresponds to a map in which galaxies are randomly distributed with random ellipticities and there is no weak lensing signals. The error bars are given by the 1σ deviations from 10 realizations. The area is the same as that simulated for the open model and the smoothing angle is $\theta_f = 1'$. The solid curve shows the prediction for a random Gaussian field.